

Influenza Situation Report

2015-2016 Seasonal Influenza Week 13

(Mar 27 2016 – Apr 2 2016)

Marion County Public Health Department, IN 2016-04-08

Please send comments or suggestions about content to epidemiology@hhcorp.org.

To subscribe to this report, email your name and email address to sjacobs@hhcorp.org, with the subject "Add to Weekly Flu Report distribution list".

Updates on Most Recent Local Activity: Week 14 (Apr 3 – Apr 9)


- The number of visits for ILI continues to decrease from the peak 4 weeks ago. The percent and count of ILI visits for the most current week (Week 14) are 1.31% and 108, respectively.
- Sixteen influenza-related deaths have been reported in Marion County this flu season.
- The Centers for Disease Control and Prevention recently reported preliminary overall influenza vaccine effectiveness (VE) of 59% this season.¹ Even though the vaccine is not 100% effective, those who are vaccinated will often have milder symptoms and fewer hospitalizations.²

Local (Week 13)

- The percent and count of ED visits with a chief complaint of ILI was 1.62 and 182 of all ED visits, respectively.
- Children age of 5-17 have the highest age-specific percent ILI (2.3%).


Emergency Department Visits for Influenza-Like Illness

Figure 1. Percent Influenza-like Illness of all Marion County Hospital ED Visits, Current Year



Data Source: ISDH ESSENCE

Figure 2. Weekly count of ILI ED Visits for Marion County Hospitals, by Age Group



Data Source: ISDH ESSENCE

Table 1. Age group specific percent of ILI ED Visits at Marion County Hospitals, by Age Group

Age Group (years)	Week 13 %ILI	Week 9-12 Average %ILI (Avg.)
<5	2.0%	6.2%
5-17	2.3%	4.5%
18-44	2.0%	2.7%
45-64	1.3%	2.0%
65+	0.4%	0.8%
Total	1.6%	2.9%

Figure 1 represents the percent of all ED visits that were categorized as ILI. The overall percent ILI for Week 13 is 1.6% (lower than last week).

ILI visit means that the patient reported fever with cough or sore throat, or used the word "flu" to describe either their symptoms or their reason for the ED visit.

Influenza Situation Report

2015-2016 Seasonal Influenza Week 13

(Mar 27 2016 – Apr 2 2016)

Marion County Public Health Department, IN 2016-04-08

Please send comments or suggestions about content to epidemiology@hhcorp.org.

To subscribe to this report, email your name and email address to siacobs@hhcorp.org, with the subject "Add to Weekly Flu Report distribution list".

Indiana (Week 13)


http://www.in.gov/isdh/files/Weekly_Influenza_Report-Week_13-2015-16.pdf

- Indiana is experiencing minimal and low influenza activity. For Week 13, ISDH reports 1.8% ILI at sentinel provider sites and 1.58% ILI ED visits. This is a decrease in sentinel surveillance and a decrease in ED surveillance compared to last week.
- Eight (8) deaths have been reported in Indiana for Week 13. The total number of mortalities in Indiana this flu season is 53.
- For Week 13 and among samples tested at the state lab: 5 specimens tested positive for 2009 A/H1N1pdm virus; 0 specimens tested positive for Influenza A/H3 seasonal virus; 0 specimens tested positive for Influenza A/H1 seasonal virus; 2 specimens tested positive for Influenza B seasonal virus; and there was 1 inconclusive sample.

National (Week 13)

<http://www.cdc.gov/flu/weekly/>

Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet
2015-16 Influenza Season Week 13 ending Apr 02, 2016



- During Week 13, the majority of states are reporting minimal and low influenza activity. In addition, during Week 13, 2.4% of outpatient visits reported through the U.S. Outpatient Influenza-like Illness Surveillance

Influenza Situation Report

2015-2016 Seasonal Influenza Week 13

(Mar 27 2016 – Apr 2 2016)

Marion County Public Health Department, IN 2016-04-08

Please send comments or suggestions about content to epidemiology@hhcorp.org.

To subscribe to this report, email your name and email address to sjacobs@hhcorp.org, with the subject "Add to Weekly Flu Report distribution list".

Network (ILINet) were due to ILI. This is a decrease from previous week (Week 12=2.9%) and is above the national baseline (2.1%).

- Seven influenza-related pediatric deaths were reported for Week 13. The season mortality is at 40.
- The most frequently identified influenza virus type reported by public health laboratories during Week 13 was influenza A, with influenza A (H1N1) viruses predominating.

Resources:

1. <http://www.cdc.gov/media/releases/2016/flu-vaccine-60-percent.html>
2. <http://www.cdc.gov/flu/about/qa/misconceptions.htm>